



Section 24(c) Special Local Need Label

**SUPPLEMENTAL LABELING FOR SPECIAL LOCAL NEED IN
BLUEBERRIES**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

Bravo Ultrex®

EPA REG. NO. 50534-201-100

EPA SLN NO. FL-040008

For control of rust diseases in blueberries after harvest

Active Ingredient: Chlorothalonil.	82.5%
Other Ingredients:	17.5%
Total	100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard restrictions and precautions on the EPA-registered Bravo Ultrex label.

This label must be in the possession of the user at the time of application.

General Information

Blueberries

After all berries are harvested, to maintain healthy leaves (thereby increasing photosynthates and subsequent plant health for the following fruiting season), apply Bravo Ultrex at 2.7 – 3.6 lbs/acre in enough water to obtain adequate coverage (normally 20-100 gallons per acre). Repeat at 10-14 day intervals.

DO NOT combine Bravo Ultrex in spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective, and noninjurious under your conditions of use. **DO NOT** combine Bravo Ultrex with Dipel , Latron AG-98 or Latron B-1956 as phytotoxicity may result from this combination when applied.

Notes: DO NOT apply more than 10.9 pounds of Bravo Ultrex (9.0 lbs.a.i.) per acre during each growing season (this includes any pre-harvest sprays).

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash waters. **DO NOT** apply when weather conditions favor drift from treated areas. **APPLY ONLY TO AREAS SPECIFIED ON LABEL.**

Section 24(c) Registrant:

GB Biosciences Corp.
PO Box 18300
Greensboro, NC 27419-8300

24(c) Sub-registrant:

Syngenta Crop Protection, Inc.
PO Box 18300
Greensboro, NC 27419-8300

EPA SLN No.:

FL-040008

Effective Date:

September 1, 2004

FL0201028AA0904

FLBravoUltrexBlueberriesSLNLbl 9-1-04

FIFRA

Section 24(c) Special Local Need Label

SUPPLEMENTAL LABELING FOR SPECIAL LOCAL NEED IN BLUEBERRIES

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

Bravo WeatherStik®

EPA REG. NO. 50534-188-100
EPA SLN NO. FL-040007

For control of rust diseases in blueberries after harvest

Active Ingredient: Chlorothalonil.	54.0%
Other Ingredients:	46.0%
Total	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard restrictions and precautions on the EPA-registered Bravo Weatherstik label.

This label must be in the possession of the user at the time of application.

General Information

Blueberries

After all berries are harvested, to maintain healthy leaves (thereby increasing photosynthates and subsequent plant health for the following fruiting season), apply Bravo Weather Stik at 3-4 pints/acre in enough water to obtain adequate coverage (normally 20-100 gallons per acre). Repeat at 10-14 day intervals.

DO NOT combine Bravo Weather Stik in spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective, and noninjurious under your conditions of use. **DO NOT** combine Bravo Weather Stik with Dipel, Latron AG-98 or Latron B-1956 as phytotoxicity may result from this combination when applied.

Notes: DO NOT apply more than 12 pints of Bravo Weather Stik (9.0 lbs.a.i.) per acre during each growing season (this includes any pre-harvest sprays).

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash waters. **DO NOT** apply when weather conditions favor drift from treated areas. **APPLY ONLY TO AREAS SPECIFIED ON LABEL.**

Section 24(c) Registrant:

GB Biosciences Corp.
PO Box 18300
Greensboro, NC 27419-8300

24(c) Sub-registrant:

Syngenta Crop Protection, Inc.
PO Box 18300
Greensboro, NC 27419-8300

EPA SLN No.:

Effective Date:

FL-040007

September 1, 2004

FL0188029AA0904

FLBravoWSBlueberriesSLNLbl 9-1-04